

THE CITY		OF BOSTON	
1787-1800		1801-1850	
Year	Population	Year	Population
1787	1,000	1801	1,500
1788	1,100	1802	1,600
1789	1,200	1803	1,700
1790	1,300	1804	1,800
1791	1,400	1805	1,900
1792	1,500	1806	2,000
1793	1,600	1807	2,100
1794	1,700	1808	2,200
1795	1,800	1809	2,300
1796	1,900	1810	2,400
1797	2,000	1811	2,500
1798	2,100	1812	2,600
1799	2,200	1813	2,700
1800	2,300	1814	2,800
		1815	2,900
		1816	3,000
		1817	3,100
		1818	3,200
		1819	3,300
		1820	3,400
		1821	3,500
		1822	3,600
		1823	3,700
		1824	3,800
		1825	3,900
		1826	4,000
		1827	4,100
		1828	4,200
		1829	4,300
		1830	4,400
		1831	4,500
		1832	4,600
		1833	4,700
		1834	4,800
		1835	4,900
		1836	5,000
		1837	5,100
		1838	5,200
		1839	5,300
		1840	5,400
		1841	5,500
		1842	5,600
		1843	5,700
		1844	5,800
		1845	5,900
		1846	6,000
		1847	6,100
		1848	6,200
		1849	6,300
		1850	6,400

a	b	c
<p>1. <i>Phragmites australis</i> (Cav.) Trin. ex Steud.</p>	<p>1. <i>Phragmites australis</i> (Cav.) Trin. ex Steud.</p>	<p>1. <i>Phragmites australis</i> (Cav.) Trin. ex Steud.</p>
<p>2. <i>Scirpus americanus</i> (L.) Stapf</p>	<p>2. <i>Scirpus americanus</i> (L.) Stapf</p>	<p>2. <i>Scirpus americanus</i> (L.) Stapf</p>
<p>3. <i>Cyperus tenuiflorus</i> (L.) Rostk Schmidt</p>	<p>3. <i>Cyperus tenuiflorus</i> (L.) Rostk Schmidt</p>	<p>3. <i>Cyperus tenuiflorus</i> (L.) Rostk Schmidt</p>
<p>4. <i>Eleocharis acicularis</i> (L.) Rostk Schmidt</p>	<p>4. <i>Eleocharis acicularis</i> (L.) Rostk Schmidt</p>	<p>4. <i>Eleocharis acicularis</i> (L.) Rostk Schmidt</p>
<p>5. <i>Eleocharis obtusa</i> (L.) Rostk Schmidt</p>	<p>5. <i>Eleocharis obtusa</i> (L.) Rostk Schmidt</p>	<p>5. <i>Eleocharis obtusa</i> (L.) Rostk Schmidt</p>
<p>6. <i>Eleocharis tenuis</i> (L.) Rostk Schmidt</p>	<p>6. <i>Eleocharis tenuis</i> (L.) Rostk Schmidt</p>	<p>6. <i>Eleocharis tenuis</i> (L.) Rostk Schmidt</p>
<p>7. <i>Eleocharis acicularis</i> (L.) Rostk Schmidt</p>	<p>7. <i>Eleocharis acicularis</i> (L.) Rostk Schmidt</p>	<p>7. <i>Eleocharis acicularis</i> (L.) Rostk Schmidt</p>





